

THE UNITED STATES OF AMERICA

<u>TO ALL TO WHOM THESE; PRESENTS SHALL, COME:</u>

AexGen Jurf Research, FIG

There has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT. THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC PRESENTAINENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR TING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PROSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

FESCUE, TALL

'Brockton'

In Testimonn Marrest, I have hereunto set my hand and caused the seal of the Plant Mariety Isotertion Office to be affixed at the City of Washington, D.C. this fourteenth day of November, in the year two thousand and eight.

Allert

Olm 3-

Commissioner Plant Variety Protection Office Agricultural Marketing Service Journal T. Schaffer

Secretary of

(See reverse for instructions and information collection burden statement)

Exhibit A:

Origin and Breeding History **Drockton* **ATF116**Tall Fescue

1.

The tall fescue (Festuca arundinacea) cultivar ATF1167 traces its parentage to the released cultivar Penn 1901. A plant selection field containing 2,500 plants of Penn 1901 was planted in the fall of 1996. In the spring/summer of 1997 the 2,500 plants were infected by stem rust incited by *Puccinia graminis*. The single plants were then rated for degree of infection. The plant selection field was flailed in early summer and the plants were rated for recovery, genetic color and crown density. One hundred plants were then selected and moved together in isolation. Following harvest in 1998 a plant selection field was established containing 1,207 plants. In the spring of 1999 the single plants were rated for dark genetic color, crown density, number of inflorescence, level of endophyte infection (Neotyphodium coenophialum) and degree of stem rust (Puccinia graminis). Forty-two clones were selected and moved to isolation blocks in the spring before anthesis. The 42 clones were harvested in bulk. In the fall of 1999, a single spaced-plant nursery was established containing 2, 200 plants. The single plants were rated for dark genetic color, crown density, number of inflorescence, level of endophyte infection (Neotyphodium coenophialum) and degree of stem rust (*Puccinia graminis*). Fifty clones were selected and moved to an isolation block in the spring before anthesis. The 50 clones were harvested in bulk.

Drought resistance has been associated with deeper root penetration for a number of turfgrass species (Burton et al., 1954; Carrow 1996). A greenhouse screening technique was developed using flexible root tubes to evaluate root extension and root length densities (Bonos et al., 2004). Clear polyethylene tubing was filled with silica sand grade No. 8 mixed evenly with 1.0 g of Scotts micro-nutrients, 2.0 g of Scotts starter fertilizer, and 4.0 g of Scotts Ploy S. The flexible tube was inserted inside a

In 2001 plants were established in rooting tubes.

PVC pipe cut to 63.5 cm, maintained at a 30 degree angle. The plants were irrigated by a drip system. Each tube was equipped with a drip emitter and received 5.9 mL of water during each irrigation cycle. The greenhouse was maintained at 25 degree C. Supplemental lighting with high pressure sodium lamps was used to maintain 12 hours of light.

One-hundred sixty-eight tubes represent a replication. This test contained three replications for a total of 504 plants. The plants were clipped six times beginning March 22, 2001 and ending April 24, 2001. The roots were harvested on April 25, 2001. The lower one foot of the flexible tube is cut and the roots are washed from the silica sand, dried and weighed. In the first replication the selection criteria was clipping average less than 0.28 g and root weight greater than 0.40 g. Six plants were selected. The second replication the selection criteria was clipping average less than 0.23 g and root weight greater than 0.50 g. Eight plants were selected. The third replication the selection criteria was clipping average less than 0.29 g and root weight greater than 0.54 g. Seven plants were selected.

The 21 plants were moved to an isolated crossing block and harvested in bulk in 2002. The population was then planted in a single spaced plant nursery in the fall of 2002. The population was replicated three times with 500 plants per replication, for a total of 1500 plants. The plants were evaluated for genetic color, freedom from disease, heading date and crown density. In the spring of 2003, 29 plants were selected and moved before flowering to an isolated crossing block. The seed was harvested in bulk and planted in a 2003 turf trial near Salem, NJ.

The turf trial was evaluated for performance and disease through the summer of Brockton?

2004. ATF116 performed well in the trial. A 1500 single plant increase block was (62:10/3/00) established in the fall. In the summer of 2005, the block was harvested in bulk and designated ATF1167, breeder seed.

References:

- 1. Burton. G.W., E. H. DeVane, and R. L. Carter. 1954. Root penetration distribution and activity in southern grasses measured by yields, drought symptoms, and P32 uptake. Agron. J. 46:229-233.
- 2. Carrow, R. N. 1996. Drought avoidance of diverse tall fescue cultivar. Crop Sci. 36:371-377.
- 3. Bonos, S.A., W. A. Meyer, D. Rush, K. Hignight. 2004. Selection for deep root production in tall fescue and perennial ryegrass. Crop Sci. 44:1770-1775.

2. Breeder Seed Maintenance:

A breeder seed multiplication was planted in isolation in 2004 in Albany, Oregon. Seed was harvested in bulk in 2005 and is maintained in cold storage. Seed propagation is limited to three generations, one each of foundation, registered, and certified.

3. Stability and Uniformity:

ATF1167 has been a stable uniform cultivar over two generations. No off-type or (be:10/3/08)
variant plants have been observed during the multiplication or reproduction. Turf plots of ATF1167 have been uniform and stable.
(be:3/10/08)

Exhibit B:

Novelty Statement of ATF1167 Tall Fescue

The following summary outlines the distinctive characteristics of ATF1167: The novelty of ATF1167 is based on the unique combination of these characteristics. ATF1167 is most similar to Rebel II, but may be differentiated by using the following criteria:

- 1) The genetic color of ATF1167 is significantly darker compared to Rebel II (tables 1A, 1B).
- 2) ATF1167 has a shorter mature plant height than Rebel II (tables 1A, 1B).
- 3) The panicle length of ATF1167 is shorter compared to Rebel II (tables 1A, 1B).
- 4) The flag leaf characteristics length, sheath length and internode length are all shorter for ATF1167 compared to Rebel II (tables 1A, 1B).
- 5) The leaf blade characteristics length and sheath length are all reduced for ATF1167 compared to Rebel II (tables 1A, 1B).
- The length of the panicle from the lower most whorl to the tip is reduced for ATF1167 compared to Rebel II (tables 2A, 2B, illus. 1).

*<ATF1167>= Brockton' (BT:5/29/08) REPRODUCE LOCALLY. Include form number and date on all reproductions.

Form Approved - OMB No. 0581-0055

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, age, disability, and where applicable, sex marital status, familial status, parental status, religion, sexual orientation, genetic infromation, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program (Not all prohibited bases apply to all programs.)

Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write\ to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6832 (TDD). USDA is an equal opportunity provider and employer.

Exhibit C

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVIC SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

OBJECTIVE DESCRIPTION OF VARIETY TALL & MEADOW FESCUES(Festuca spp.)

NAME OF APPLI	CANT(S)		ITEMP	ORARY DESIGNATIO	N I		VAR	IE I Y	NAM	E			
NexGe	en Turf Research, I	LLG _{*:j}		ATF1167	İ	B	rock	ton) (bt.)	19/3/	¹ '08)		
ADDRESS (Street	et and No., or RD No.,	City, State, ZIP Cod	de, and Country)		ļ	FOR	OFFIC	IÁL U	SE O	NLY			
	Columbus ST S.E. , QR, USA				.]	PVP Ø (1	210	FR ₀	8	0 (0 (8 (4
PLEASE READ A	ALL INSTRUCTIONS	CAREFULLY:											,
Place the appropriat	te number that describes istics described including	the varietal characteristi	nts, should represent th	coxes below. Use leading a nose that are typical for the ultural conditions must be	variety. Meas	ured dat	a should	be fo	r SPAC	CEDF	LIAN	TS. GI	ve addition:
1. SPECIES: (Wi	th comparison varietie	es, use varieties with	in the species of the	application variety)									
61 = F. arur	ndinacea (Tall)			<u>Turf Types</u>									
	1 = Kentucky 31	2 = Rebel	3 = Olympic	4 = Bonanza	5 = Arid		6 = F	Rebel	11				
	7 = Shortstop	8 = Silverado	9 = Rebel Jr.	10 = Mini Mustang	11 = Crewo	cut	12 =	Bons	sai				
			Forage	<u>Types</u>									
	20 = Kentucky 31	21 = Martin	22 = Forager	23 = Mozark									
	24 = Kenhy	25 = AU Triumph	26 = Fawn	27 = Cajun									
2 = F. prate	ensis (Meadow)												
	30 = Admira	31 = Beaumont	32 = Comtessa	33 = Ensign	34 = Trade	ır							
2. CYTOLOGY:													
2N=42 Ch	romosome Number												
3. ADAPTATION:	(0 = Not Tested; 1 =	Not Adapted; 2 = Ad	apted)										
0T	ransition Zone	2West	2_ Northeast	Other (Spec	ify):				_				
4. MATURITY: (I	Date First Headed, 10	0% of Panicle Emerg	ence)										
5 Maturity CI	ass 1 = Verv	reariy () 2 = AU	Friumph 3 = Ear	rly (Fawn,) 4 = Kenh	y 5=Me	dium (R	lebel)						
	6 = Bona		(Silverado) 8 = Ver	rv late	-								
		67 Days after April 1											
	Avg. No. days to h												
		<u> </u>		_									

4. MATURITY: (continued)	
Days earlier than	#200800081
Maturity same as6 Comparison Varie	etv
Days later than	•
5. MATURE PLANT HEIGHT cm: (Average of 100 culms from crown to top of panicle, if panicle is nodding, straighten)	INTERNODE LENGTH CM: (First internode subtending the flag leaf)
<u>90.80</u> cm Height	14.40 cm Internodel_ength
20.20 cm Shorter than _6_	5.23 cm Shorter than6
Height same as Comparison Variety	Length same as Comparison Variety
cm Taller than	cm Longer than
HEIGHT AT EAR EMERGENCE cm: (Flag leaf height from crown to flag leaf	node)
41.10 cm Height	
15.40 cm Shorter than _6_	
Height same as Comparison Variety	
cm Taller than	
6. GROWTH HABIT: (Mature Plants)	
9 6 = Prostrate () 3 = Semiprostrate (6)	5 = Horizontal ()
7 = Semierect (Rebel) 9 = Erect (Mini Mustang)	
7. RHIZOMES: (Psuedo):	
mm Length6_1 = Absent () 2 = Rare (Rebel)	3 = Common ()
8. LEAF BLADE: (Tiller leaves/ turf color)	
	green () 5 = Green ()
7 = Medium dark green () 9 = Very dark gree	
4 Specify Rating of Comparison Variety	
1 Anthocyanin: 1 = Absent (6) 9 = Present ()	
1 Basal Hairs: 1 = Absent (6) 9 = Present ()	
) 9 = Rough (·)
_6_Width Class: 1 = Very coarse () 3 = Coarse (6)	5 = Medium ()
7 = Fine () 9 = Very Fine (
	, LER LEAF WIDTH MM:
	67_ mm Tiller Leaf Width
6.73_ cm Shorter than6	mm Narrower than
Length same as Comparison Variety	Width same as _6_ Comparison Variety
cm Taller than	mm Longer than

8. LEAF BLADE: (Continued)		# 2	200800081
FLAG LEAF LENGTH CM:		FLAG LEAF WIDTH MM:	
39.17 cm Flag Leaf Length		7.33 mm Flag Leaf Width	
8.86 cm Shorter than _6_		mm Narrower than	-)
Length same as	Comparison Variety	Width same as _6	Comparison Variety
cm Longer than J		mm Wider than	-)
9. LEAF SHEATH: (Basal Portion)			
Anthocyanin (seedling): 1 = Absent (KY	′-31) 9 = Prese	ent ()	
9 Auricle Hairiness: 1 = Absent () 9 = Prese	ent (6) 97% See table 4	
10. PANICLE: (At seed maturity except where noted.)			
7 Shape: 1 = Narrow-tapering ()	5 = Ovate ()	7 = Oblong ()	9 = Other (specify)
5 Type: 1 = Compact (appressed)) 5 = Intermediate () 7 = Open ()	9 = Other (specify)
9 Orientation: 1 = Nodding (9 = Erect ()	
1 Branch Pubescence: 1 = Glabrous (6)	9 = Pubescent ()	
	'ellowish Green Purplish	2 = Green 3 = Bluish Green 5 = Reddish 6= Other (Specify)
	'ellowish Green Purplish	2 = Green 3 = Bluish Green 5 = Reddish 6= Other (Specify)
72.03 cm Panicle Length (from base to tip, i	if nodding, straighten; aft	er anthesis)	
11.39 cm Shorter than _6_			
Length same as	Comparison Variety		
cm Longer than J			
11. SEED: (With Lemma & Pelea)			
*_2877 mg per 1000 seeds			·
mg Less than			
Weight same as	Comparison Variety		
PALEA: (Keels or Margins) _3_ Hairs: 1 = A	ubsent()	5 = Short (Missouri 96)	9 = Long ()
LEMMA: _1_ Hairs: 1 = A	bsent (Kenhy)	5 = Several ()	9 = Many (Missouri 96)
_6.53 mm Lemma Length (Mature)		1.20 mm Lemma Width	
mm Shorter than		mm Narrower than	-)
Length same as _6_ Co	mparison Variety	Width same as6	Comparison Variety
mm Longer than		mm Wider than	- J

11. SEED: (conti	inued)					#2	0 0	8 0	0 0	8	1
AWNS	S: _9_AWNS:	1 = Absent ()	9 = Present (Fa	alcon)	_100_% Plants w			- +			2
1.27	_ mm Awn length (Of th	nose present.)									
<u></u> -	mm Shorter than	_)									
!	Length same as	_6_	mparison Variet	y							
	mm Longer than	_)									
12. DISEASE, IN	ISECT, AND NEMATO	DE REACTION: (0=	= Not Tested 1= L	east Resistan	: 9= Most Resistar	nt)					
0 M	elting-out (<i>Drechslera</i> ,	poae)		_0_ Blind	l Seed (Gloeotinia	temulenta)					
0 Le	eaf Spot (D. siccans)			_0_ Dolla	ar Spot (<i>Lanzia, Mo</i>	ollerdiscus spp	.)				
0 Ne	et Blotch (D. dictyoides	;)		_0_ Sten	n Rust (<i>Puccinia gr</i>	ramminis)					
0 Br	own Patch (<i>Rhizoctoni</i>	ia solani)		_0_ T. BI	ight (<i>Typhula incar</i>	rnate)					
0 C.	Leaf Spot (Cercospor	a fectucae)		_0_ Pyth	ium Blight (<i>Pythiun</i>	n spp.)					
0 Pi	nk Snow Mold (<i>Gerlac</i> i	hia nivalis)		_0_ Pow	dery Mildew (<i>Erysi</i> į	phe gramminis)				
0 Sil	ver Top (F. tricinctum,	F. roseum)		_0_ Crow	n Rust (<i>Puccinia d</i>	coronata)					
Ot	her Disease										
Ot	her Insect										
Ot	her Nematode	- 13 d 18 to - 2 to									
13. ENVIRONME	NTAL STRESS:			· · · · · · · · · · · · · · · · · · ·							
5 Dr	ought Stress	1 = Susceptible () 5 = Te	olerant (6)	9 = Res	sistant ()					
5 Sh	ade Stress	1 = Susceptible () 5 = To	olerant (6)	9 = Res	sistant ()					
5 Wi	nter Stress	1 = Susceptible () 5 = To	olerant (6)	9 = Res	sistant ()					
14. GIVE VARIET esemblance with	TY OR VARIETIES THAT the following scale:	AT MOST CLOSEL'	Y RESEMBLE TH	E APPLICATION	ON VARIETY. For	the following o	haracter	istics, i	ndicate	the d	egree of
I = Application va	ariety is less than comp	parison variety.	2 = Same as	3 = More	than, Better, Great	ter, Darker, etc					
Character	Varieties	Rating			Character	Varieties		Rating			
_eaf Width	REBEL II	1		and a second of	Leaf Color	REBEL II	. —	3			
Panicle Color	REBEL II	2			Panicle Shape	REBEL II		2			

Character	Varieties	Rating	Character	Varieties	Rating
Leaf Width	REBEL II	1	Leaf Color	REBEL II	3
Panicle Color	REBEL II	2	Panicle Shape	REBEL II	2
Seed Size	REBEL II	1	Cold Injury	REBEL II	3
Winter Color	REBEL II	3	Heat	REBEL II	3
Disease	REBEL II	3			

15. EXPERIMENTAL: Give a brief summary of the experimental design utilized to collect the data used on this form. Cultural conditions, number of plants measured and plant spacing must be specified.

A morphological nursery designated 05PVPFA1 was established in September of 2005, in Albany, Oregon. Experimental design consisted of 9 entries; 3 replications per entry; 20 plants per replication; for a total of 60 plants per entry. Crewcut, Rebel II, and KY-31were used as standards. Plants were established on 2.5 foot centers with a skip row between replications and between entries.

The nursery received 30 pounds of nitrogen per acre rate following establishment and 50 pounds of nitrogen per acre per year in 2006 and 2007. The fertilizer source was 15-15-15 and was applied as a split application with 1/2 applied in the spring and 1/2 in the fall. The nursery was sprayed twice each spring, 3 weeks between applications, with Tilt (2 oz/acre rate), to prevent stem rust. One pound of Karmex per acre rate was applied during late summer to prevent emergence of volunteer seedlings.

Data was analyzed using analysis of variance for a randomized complete block design. Means were calculated for each replication and then analyzed.

16. ADDITIONAL DESCRIPTION: (Use additional sheets as required).

Describe all characteristics that cannot be adequately described in the form above in Exhibit D. Varieties used for comparison should be used as may be appropriate, such as for disease reactions. Append all comparative trial and evaluation data, including measured characters, environmental, and disease tests results. Providing such information may aid in conducting a more thorough review of the applicants claims of distinctness.

#200800081

Exhibit D:

Additional Description Brock for? ATFTI6>Tall Fescue (bt:10/3/08)

Brockton'
ATF1167 is an improved turf-type tall fescue. It has a shorter mature plant height (tables 1A, 1B) than previously released tall fescue cultivars, such as KY-31 and Rebel II. ATF1167 has a medium maturity with an anthesis date earlier than Crewcut (tables 1A, 1B). ATF1167 exhibits a darker genetic color compared to KY-31 and Rebel II (tables 1A, 1B). The panicle length of ATF1167 is shorter compared to KY-31 and Rebel II (tables 1A, 1B). The flag leaf characteristics; length, sheath length and internode length are all shorter for ATF1167 than KY-31 and Rebel II (tables 1A, 1B). The flag leaf height, and internode length of ATF1167 is reduced compared to Crewcut (tables 1A, 1B). The leaf blade characteristics of blade length and height are shorter for ATF1167 than KY-31, Rebel II and Crewcut (tables 1A, 1B). ATF1167 has a shorter panicle length from the lower most whorl to the tip, compared to KY-31, Rebel II, and Crewcut (tables 2A, 2B). The whorl characteristic, length of longest branch is shorter for ATF1167 compared to KY-31 (tables 2A, 2B, illus. 1). The distance between the two most lower whorls is less for ATF1167 compared to Crewcut and KY-31 (tables 2A, 2B). The number of spikelets on the longest branch of the lower most whorl is greater for ATF1167 compared to Rebel II and KY-31 (tables 2A, 2B). ATF1167 has fewer plants with purple pigmentation in the panicles than Rebel II (tables 3A, 3B). ATF1167 has a more erect growth habit compared to KY-31 (tables $\frac{2}{3}$ A, $\frac{2}{3}$ B). The presence of only one branch on the lower whorl is more frequent in ATF1167 than KY-31 (tables 3A, 3B). The expression of two branches on the lower most whorl is less for ATF1167 compared to Rebel II and KY-31 (tables 3A, 3B). ATF1167 expresses fewer plants with dark pigmentation at the nodes compared to KY-31 and Rebel II (tables 4A, 4B). ATF1167 has a higher seed weight per 1,000 seeds compared to Rebel II and Crewcut, but less than KY-31 (tables 4A, 4B).

Table 1A	<	٠.			200	2006 Morphological Data	oloda	gical	Data						
Cultivar	Genetic	Heading	Anthesis	Mature	置	Plant Panicle	Flag	Flag	_	Flag	Flag	Leaf	Leaf	Leaf	Leaf
	Color	Date	Date	Plant	Width	rength	Leaf	Leaf	Leaf	Leaf	Leaf	Blade	Blade	Blade	Sheath
***************************************	(1-9 Scale	ใ-9 Scale (days after	(days after	Height	(cm)	(cm)	Length Width	Width	Height	Sheath	Height Sheath Internode Length	Length	Width		Length
	9≃Best)	March 1)	April 1)	(cm)			(cm)	(mm) (cm)		Length	Length Length	(cm)	(mm) (cm)		(cm)
										(cm)	(cm)				
* ATF1167	6.00	35.67	55.33	90.80	24.03	90.80 24.03 72.03	39.17 7.33	7.33	41.10	41.10 24.53	14.40	33.80	9.67	16.83	13.23
CE3	6.00	30.67	54.33	95.20	95.20 24.00	75.87	36.37	6.00	42.67	23.57	14.40	30.07	8.33	17.30	12.47
RP1	6.33	34.00	56.33	89.03	89.03 23.50	69.10	33.47	5.33	42.37	22.20	15.23	28.37	8.00	17.17	12.03
ATF1203	7.00	37.33	58.67	88.63	22.93	88.63 22.93 70.07	37.70	6.67	42.50	24.03	13.87	32.97	32.97 10.00 17.50	17.50	13.07
Crewcut	6.00	38.33	60.67	97.37	97.37 24.13	73.00	39.07	6.67	48.07	24.40	18.07	36.57	9.67	9.67 20.13	14.10
Rebel II	4.67	35.33	59.33	111.00 24.87	24.87	83.43	48.03	7.33	56.50	29.90	19.63	40.53	40.53 10.33 23.33	23.33	16.50
KY-31	3.67	33.00	60.00	130.33	21.20	130.33 21.20 92.87	51.87	7.67	69.73	35.40	23.30	45.20	45.20 11.00 33.37	33.37	20.07
LSD (0.05)		1.72	1.82	5.75	3.56	4.92	2.38	1.12	3.55	2.08	1.44	1.93	0.65	2.01	0.87
S	7.03	3.38	2.16	3.94	3.94 10.41	4.41 4.01 11.49 4.97	4.01	11.49	4.97	5.43	5.83	3.75	3.75 4.65 6.63	6.63	4.14

Cultivar under evaluation
 Significant difference over two locations one year.
 Significant difference over one location one year.
 Measurements taken in Albany, Oregon
 3 reps; 20 plants/rep = 60 data points

Table 1B	~				20(2007 Morphological Data	olodd	gical	Data						
	Genetic	Heading	Anthesis	Mature	Plant	Mature Plant Panicle	Flag	Flag	_	Flag	Flag	Leaf	Leaf	Leaf	Leaf
_	Color	Date	Date	Plant	Width	Width Length	Leaf	Leaf	Leaf	Leaf	Leaf	Blade	Blade	Blade	Sheath
_	(1-9 Scale days		after (days after Height	Height	(cm) (cm)	(GJ)	Length	Width	Height	Sheath	Height Sheath Internode Length Width	Length		Height	Length
	9=Best)	March 1)	1 1) April 1)	(GJJ)			(GII)	(mm) (cm)		Length	Length	(cm)	(mm)	(cm)	(cm)
										(cm)	(cm)				
\blacksquare	6.33	53.67	26.00	120.37 28.43	28.43	1 1	85.90 44.23	6.00	61.13	27.63	24.67	40.37	7.33	40.37 7.33 25.10	15.97
	6.33	20.00	54.67	113.03	28.80	113.03 28.80 78.40 42.30		5.00	59.63	25.17	23.47	37.83	5.67	25.90	15.00
	6.00	53.67	54.67	111.13	28.43	111.13 28.43 77.57	39.77	5.33	59.70	25.47	24.33	34.77	5.33	24.50	14.77
г	6.33	55.00	52.33	122.07	28.83	122.07 28.83 85.13	44.47	6.00	64.57	28.27	24.90	40.97	7.00	28.03	16.73
Г	5.67	58.00	29.00	125.40	30.23	125.40 30.23 86.47	44.57	00.9	66.67	27.10	26.93	44.10	7.33	29.50	17.03
	4.33	55.00	57.33	134.83	30.13	134.83 30.13 92.70	53.20	6.67	74.57	32.90	28.73	50.10	50.10 7.67	31.97	20.03
Г	4.00	52.00	55.67	155.17	28.70	155.17 28.70 97.10	57.43	7.00	7.00 96.27	38.30	29.57	97.60	9.67	52.33	24.37
SD (0.05)	99.0	1.61	1.44	5.64	2.16	5.57	3.45	0.86	3.87	1.61	1.41	3.65	0.87	3.28	1.08
	8.15	2.05	1.76	3.08	3.08 5.10	4.45	60'9	9.85	9.85 3.86	3.77	3.72	5.75	8.37	7.27	4.18

Example 2 Contiver under evaluation
Significant difference over two locations one year.
Significant difference over one location one year.
Measurements taken in Albany, Oregon
3 reps; 20 plants/rep = 60 data points

*CATF 1167>= 'Brocktori' (BT: 10/3/2008)

Table 2A	V				2006	Labora	2006 Laboratory Morphological Data	hological	Data				
Cultivar	Lemma	Lemma Lemma	Lemma	Palea	Palea	Glume	Glume Length of	Spikelets		Spikelet	Spikelet Length of	Distance	Number of
	Length	Width	Awn	Length Width	Width	Length	Length Panicle from per Panicle per	per Panicle		Length			Spikelets on
	(mm)	(mm)	Length	(mm) (mm)		(mm)	Lower Most		kelet	(mm)		S	the Longest
*****			(mm)				Whorl to Tip				Lower Whorl	Whorls	Branch
							(mm)				(mm)	(mm)	Lower Whorl
* ATF1167	6.53	1.20	1.27	6.47	1.13	5.07	235.43	103.33	6.67	12.33	112.77	60.77	21.33
CE3	6.57	1.27	1.37	6.30	1.13	4.90	225.33	77.00	7.33	12.50	104.93	57.33	14.00
RP1	6.40	1.30	1.23	6.23	1.13	4.70	207.40	76.00	7.33	12.27	94.73	52.53	14.00
ATF1203	6.70	1.27	1.20	6.37	1.13	4.77	227.87	106.67	6.33	11.83	103.47	54.97	20.33
Crewcut	7.00	1.47	1.67	6.50	1.23	5.10	264.50	109.33	7.33	12.90	126.17	67.67	19.67
Rebel II	7.40	1.43	1.73	6.70	1.20	5.23	279.93	112.33	6.67	12.37	125.53	69.23	18.67
KY-31	7.80	1.40	1.30	7.43	1.27	5.63	324.87	120.33	8.00	14.07	138.57	80.40	17.00
LSD (0.05)	0.48	0.23	0.31	0.32	0.12	0.35	15.62	19.37	08.0	1.04	10.52	3.03	2.39
<u>ک</u>	4.74	11.92	15.34	3.34	6.78	4.77	4.26	13.21	7.74	5.66	6.28	3.29	9.20

Euitivar under evaluation
 Significant difference over two locations one year.
 Significant difference over one location one year.
 Measurements taken in Albany, Oregon
 3 reps; 20 plants/rep = 60 data points

Table 2B	മ			- 1	2007	Labora	2007 Laboratory Morphological Data	Shological	Data				
Cultivar	Lemma	emma Lemma	Lemma Palea		Palea	Glume	Glume Length of	Spikelets	Florets	Spikelet	Spikelet Length of	Distance	Number of
	Length	Width	Awn	Length Width	Width	Length	Length Panicle from per Panicle per	per Panicle	per	Length		Between	Spikelets on
	(mm)	(mm)	Length	(mm)	(mm)	(mm)	Lower Most		Spikelet	(mm)	Branch	Lower Most	the Longest
			(mm)				Whorl to Tip				Lower Whorl	Whorls	Branch
							(mm)				(mm)	(mm)	Lower Whorl
ATF1167	6.43	1.27	1.27	6.53	1.17	5.13	279.73	109.00	5.67	10.83	113.50	72.03	19.00
CE3	6.07	1.27	1.27	6.47	1.23	4.90	250.63	80.00	5.67	11.23	93.07	60.30	12.67
RP1	6.33	1.27	1.07	6.27	1.13	4.80	233.60	77.00	5.33	10.80	91.30	56.90	12.67
ATF1203	6.47	1.30	1.33	6.63	1.23	5.17	284.70	104.00	5.33	10.93	112.53	69.30	19.00
Crewcut	6.53	1.20	1.37	6.77	1.27	5.20	299.33	95.00	6.33	12.07	125.70	77.43	18.00
Rebel II	6.80	1.30	1.40	7.03	1.23	5.07	306.97	106.67	6.00	11.90	119.70	75.57	16.67
KY-31	7.00	1.33	0.93	7.13	1.20	5.47	350.87	124.00	5.67	12.40	129.13	86.57	16.33
(CSD (0.05)	0.54	0.16	0.43	0.33	0.17	0.32	17.31	7.96	0.71	0.87	12.65	4.97	1.60
ે ટ	5.68	8.46	23.69	3.34	69.6	4.31	4.15	5.51	8.54	5.20	7.76	4.80	6.75

Equitivar under evaluation

Significant difference over two locations one year.


Significant difference over one location one year.

Measurements taken in Albany, Oregon

3 reps; 20 plants/fep = 60 data points

* CATHIGT>= Brackton (BT:10/3/08)

Panicle Type Inflorescence



15.

Table 3A	Υ			2006	2006 Addition	al Morpl	nological	l Measu	ional Morphological Measurements of the Panicle	of the F	Panicle				
Cultivar	Anther	Panicle	Lemma		Panicle	Panicle	Panicle	Panicle	Panicle	Panicle Panicle	Panicle	Panicle	Panicle	Panicle Panicle Panicle	Panicle
	<u>ö</u> <u>Co</u>	io O	Awn	Color	Orientation Shape	Shape	Shape	Shape	Type	Type	Type	Branch	Branch	Branch Branch	Branch
	% Purple	% Purple	% Present	% Purple	% Nodding	% Narrow	% Oblong	% Ovate	% Purple % Purple % Present % Purple % Nodding % Narrow % Oblong % Ovate % Compact % Inter- % Open	% Inter-	% Open	Lower	Lower	Lower	Pubescence
										mediate		Whorl	Whorl	Whorl	% Present
												No. branches=1	=2	<u>۸</u>	
★ ATF1167	2	8	100	2	5	0	92	8	0	92	8	15	77	8	0
CE3	2	2	100	3	12	10	85	5	9	85	5	12	89	8	c
RP1	10	7	100	0	7	12	83	ည	12	83	ည	17	76	_	0
ATF1203	သ	10	100	ဗ	ß	9	82	12	9	85	12	27	8	13	0
Crewcut	2	10	9	က	ည	0	83	17	0	ဆ	1		88	_	0
Rebel II	က	32	100	ဗ	23	2	13	82	2	43	88	12	88	6	ဌ
KY-31	7	88	6	0	18	0	20	08	0	20	8	2	66	9	7
														•	-

■ Cultivar under evaluation Measurements taken in Albany, Oregon 3 reps; 20 plants/rep = 60 data points

l able 3B	38			2007	Addition	al Morpt	nological	l Measu	2007 Additional Morphological Measurements of the Panicle	of the F	anicle					
Cultivar	Anther Color % Purple	Panicle Color % Purple	Lemma Awn % Present	Glume Color % Purple	Anther Panicle Lemma Glume Panicle Panicle Color Awn Color Orientation Shape % Purple % Purple % Nodding % Narrow	Panicle Shape % Narrow	Panicle Shape % Oblong	Panicle Shape % Ovate	Panicle Panicle Shape Shape Shape Shape Type % Oblong % Ovate % Compact	Panicle Panicle Pranicle Type Type Branch % Open Lower mediate Whorl	Panicle Type % Open	Panicle Branch Lower Whorl No. branches=1	Panicle Panicle Panicle Branch Branch Lower Lower Pubeso Whorl Whorl % Press	Panicle Branch Lower Whorl	Panicle Branch Pubescence % Present	
ATF1167	10	∞	100	0	2	24	89	18	24	58	18	18	63	19	0	т
CE3	£	15	100	0	2	52	42	9	52	42	9	13	- 22	9	_	1
RP1	12	3	100	0	က	33	09	11	33	20	17	13	11/	9	0	Ŧ
ATF1203	8	7	100	0	2	23	47	30	23	47	30	28	28	4	0	г
Crewcut	8	12	100	0	6 0	20	48	32	20	48	32	18	25	25	0	r-
Rebel II	7	12	100	0	28	45	37	18	45	37	18	13	22	12	0	П
KY-31	8	10	100	0	32	18	25	30	18	52	30	2	2.2	21	-	_
																7

Cultivar under evaluation
 Measurements taken in Atbany, Oregon
 reps; 20 plants/rep = 60 data points

* (ATFII 677="Brocktor"/ (87810/3/2008)

Table 4A	A	:	2006 Ad	2006 Additional Morphological N	ological A	Measurements	ents							
Cultivar	Anthocyanin Present in the Leaf Blade % Purple	Leaf Blade Margin Roughness to the Touch % Smooth	Anthocyanin Leaf Blade Margin Leaf Blade Margin Present in the Roughness to the Roughness to the Leaf Blade Touch Touch % Smooth % Semi-Rough % Rough	Leaf Blade Margin Roughness to the Touch % Rough	Leaf Blade Margin Hairs % Present	Leaf Sheath Auricle Hairs % Present		Lemma Palea Node Hairs Color % Present % Distinct	Palea Hairs % Present				Growth Habit at Anthesis	Growth Habit at Anthesis
ATE1167		6							1		Seeds	% Prostrate	% Semi-erect	% Frect
2012		53	49	28	97	97	0	89	100	13	2877	6	38	53
E3	0	22	52	56	9	96	0	78	8	~	2187		2	3 8
5	0	23	55	22	ä	8	,			,		,	-	8
A TEL 2000				77	3	8	3	70	3	5	1914	0	φ	8
202		71	46	42	83	95	0	77	100	ო	2903	4	\$	62
TOMOUT Crowcut		15	53	32	100	9	0	99	180	15	2152	6	34	99
Rebell	0	35	45	20	06	90	0	92	100	42	2434	15	34	25
KY-31	0	44	34	22	8	85	0	75	100	æ	3104	8	ę	,

3 reps; 20 plants/rep = 60 data points

Table 4B	ā		2007 Additic	nal	Nological	Morphological Measurements	nents							
Cultivar	Anthocyanin Present in the Leaf Blade	Leaf Blade Margin Roughness to the Touch	Anthocyanin Leaf Blade Margin Leaf Blade Margin Leaf Blade Margin Present in the Roughness to the Roughness to the Leaf Blade Touch Touch	Leaf Blade Margin Roughness to the Touch	Leaf Blade Margin Hairs	Leaf Sheath Rhizomes Auricle % Present Hairs	ent	Lemma Hairs % Present	Palea Node Hairs Color % Present % Distinct		Seed Growth Weight Habit at	Growth Habit at	Growth Habit at	Growth Habit at
	% Purple	% Smooth	-≟I	ğ	sent	% Present					seeds)	% Prostrate	% Prostrate % Semi-erect % Erect	% Erect
A1F1167	٥	65	27	8	92	100	0	88	100	2	2853	0	33	67
CE3	0	82	18	0	82	100	0	85	199	7	2177	c	23	77
RP1	0	62	32	မ	87	100	0	92	100	_	1080	c	200	7.7
ATF1203	0	29	25	80	92	98	c	8	20	, c	2048		36	-
Crewcut	0	53	37	10	26	98	0	82	100	,	2108	0 4	386	, g
Rebel II	0	90	37	13	06	95	0	93	9	12	2498	- ~	48	3 6
KY-31	0	75	10	15	83	87	С	69	100	17	3000	·	4	3 \$
Cultivar un	Cultivar under evaluation							,	3		3000	7	20	?

■ Cultivar under evaluation Measurements taken in Albany, Oregon 3 reps: 20 plants/rep = 60 data points

K-ATF1167>='Brockton'

REPRODUCE LOCALLY. Include form number and edition date on al	I reproductions. Fo	ORM APPROVED - OMB No. 0581-0055		
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E	Application is required in order to dete certificate is to be issued (7 U.S.C. 24 confidential until the certificate is issued	21). The information is held		
STATEMENT OF THE BASIS OF OWNERSHIP				
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME		
NexGen Turf Research, LLC	OR EXPERIMENTAL NUMBER ATF1167	Bracklon?		
		(bt:19/3/08)		
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZiP, and Country)	5. TELEPHONE (Include area code)	6. FAX (Include area code)		
33725 Columbus St. S. E. Albany, OR	(541) 967-8923	(541) 967-8223		
97322	7. PVPO NUMBER	The Management of the Control of the		
71752				
8. Does the applicant own all rights to the variety? Mark an "X" in the appropriate block. If no, please explain. YES NO				
9. Is the applicant (individual or company) a U.S. national or a U.S. b				
10. is the applicant the original owner? YES	NO If no, please answer <u>one</u> o	of the following:		
a. If the original rights to variety were owned by individual(s), is (are) the original owner(s) a U.S. National(s)? YES NO If no, give name of country				
b. If the original rights to variety were owned by a company(ies), is (are) the original owner(s) a U.S. based company? YES NO If no, give name of country 11. Additional explanation on ownership (Trace ownership from original breeder to current owner. Use the reverse for extra space if needed):				
PLEASE NOTE:				
Plant variety protection can only be afforded to the owners (not licens	sees) who meet the following criteria:			
 If the rights to the variety are owned by the original breeder, that penational of a country which affords similar protection to nationals of 				
If the rights to the variety are owned by the company which employ nationals of a UPOV member country, or owned by nationals of a genus and species.				
3. If the applicant is an owner who is not the original owner, both the $\boldsymbol{\theta}$	original owner and the applicant must me	eet one of the above criteria.		
The original breeder/owner may be the individual or company who dis Act for definitions.	rected the final breeding. See Section 4	I(a)(2) of the Plant Variety Protection		
According to the Papenwork Reduction Act of 1995, an agency may not conduct or sponsor, control number. The valid OMB control number for this information collection is 0581-0055, including the time for reviewing the instructions, searching existing data sources, gathering a	The time required to complete this information collect	on is estimated to average 0.1 hour per response,		
The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and a marital or family status, political beliefs, parental status, or protected genetic information. (N communication of program information (Braille, large print, audiotape, etc.) should contact U.	ot all prohibited bases apply to all programs.) Person:	s with disabilities who require alternative means for		

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.

REPRODUCE LOCALLY. Include form number and date on all reproductions.

Form Approved OMB NO 0581-0056

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 5 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, Jarge print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE **SCIENCE AND TECHNOLOGY** PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

EXHIBIT F DECLARATION REGARDING DEPOSIT

	DECEARATION REGARDING DEFOSIT	
NAME OF OWNER (S)	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country)	TEMPORARY OR EXPERIMENTAL DESIGNATION
NexGen Turf Research, LLC	33725 Columbus St. SE. Albany, OR	ATF1167
	97322 USA	VARIETY NAME BYOCKLON'S CHE MINES
NAME OF OWNER REPRESENTATIVE (S)	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country)	FOR OFFICIAL USE ONLY
Kenneth Hignight	33725 Columbus St. SE, Albany, OR 97322 USA	PVPO NUMBER 2 0 0 8 0 0 0 8 1

I do hereby declare that during the life of the certificate a viable sample of propagating material of the subject variety will be deposited, and replenished as needed periodically, in a public repository in the United States in accordance with the regulations established by the Plant Variety Protection Office.

Kennett / Hegrifla

1-17-08